

**A SHORT PROOF OF $w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = 0$ FOR ALL n
AND A GRAPH COLOURING THEOREM BY
BABSON AND KOZLOV**

CARSTEN SCHULTZ

ABSTRACT. We show that the n -th power of the first Stiefel-Whitney class of the $\mathbb{Z}_2$ -operation on the graph complex $\text{Hom}(C_{2r+1}, K_{n+2})$ is zero, confirming a conjecture by Babson and Kozlov. This proves the strong form of their graph colouring theorem, which they had only proven for odd n . Our proof is also considerably simpler than their proof of the weak form of the theorem, which is also known as the Lovász conjecture.

1. INTRODUCTION

A homomorphism from a graph G to a graph H is a function $f: V(G) \rightarrow V(H)$ between their vertex sets that respects the edge relation, i.e. such that $\{f(u), f(v)\} \in E(H)$ whenever $\{u, v\} \in E(G)$. If the vertices of H are identified with the vertices of the simplex $\Delta^{|V(H)|-1}$, then functions $V(G) \rightarrow V(H)$ correspond to vertices of the cell complex $\prod_{v \in V(G)} \Delta^{|V(H)|-1}$. The subcomplex consisting of all cells all of which vertices correspond to homomorphisms is denoted by $\text{Hom}(G, H)$. Hom is a functor, for its basic properties, as well as background on the results mentioned in this introduction, we refer the reader to [BK06a] and [Koz06a]. We will however try to be as self-contained as possible in the proof of our main result, Theorem 1.3.

The main result of [BK06b] is the following theorem. C_s denotes the circular graph with s edges.

1.1. Theorem. *Let G be a graph, $r \geq 1$ and $n \geq 0$. If $\text{Hom}(C_{2r+1}, G)$ is $(n-1)$ -connected, then G is not $(n+2)$ -colourable.*

This had been conjectured by Lovász and is analogous to the following theorem that represented his novel approach to graph colourings in his proof of Kneser's conjecture [Lov78].

1.2. Theorem. *Let G be a graph and $n \geq -1$. If $\text{Hom}(K_2, G)$ is n -connected, then G is not $(n+2)$ -colourable.*

Lovász did originally not use the complex $\text{Hom}(K_2, G)$, but another graph complex, the neighbourhood complex, which is homotopy equivalent to it.

The proofs of both theorems rely on the study of a free $\mathbb{Z}_2$ -action on the space $\text{Hom}(C_{2r+1}, G)$ or $\text{Hom}(K_2, G)$ respectively, which is induced by the nontrivial

This research was supported by the Deutsche Forschungsgemeinschaft within the European graduate program "Combinatorics, Geometry, and Computation" (No. GRK 588/2).

automorphism of K_2 respectively an automorphism of C_{2r+1} flipping one edge. We will always regard these spaces as equipped with these actions. Similarly, when a $\mathbb{Z}_2$ -action on a sphere is mentioned, this will always be given by the antipodal map unless noted otherwise.

The basic idea of the proofs of both theorems is the same. An $(n+2)$ -colouring of G is just a graph homomorphism $G \rightarrow K_{n+2}$ and induces a map $\text{Hom}(H, G) \rightarrow \text{Hom}(H, K_{n+2})$, which, by functoriality of Hom with respect to its first argument, is a $\mathbb{Z}_2$ -map for $H = K_2$ or $H = C_{2r+1}$. If $\text{Hom}(H, G)$ is $(k-1)$ -connected, then a $\mathbb{Z}_2$ -map $\mathbb{S}^k \rightarrow \text{Hom}(H, G)$ and hence $\mathbb{S}^k \rightarrow \text{Hom}(H, K_{n+2})$ exists. A study of the space $\text{Hom}(H, K_{n+2})$ then reveals conditions on r and n under which such a map does not exist. In the case of Theorem 1.2 this amounts to the realisation that $\text{Hom}(K_2, K_{n+2}) \approx_{\mathbb{Z}_2} \mathbb{S}^n$ (see Remark 2.5) and an invocation of the Borsuk-Ulam Theorem.

Babson's and Kozlov's proof of Theorem 1.1 uses different arguments for odd and for even r ; both rely on elaborate cohomology calculations for the cell complex $\text{Hom}(C_{2r+1}, K_{n+2})$. For even n they show that a $\mathbb{Z}_2$ -map $\text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow \mathbb{S}^n$ exists such that the image of a generator of $H^n(\mathbb{S}^n)$ in $H^n(\text{Hom}(C_{2r+1}, K_{n+2}))$ under the induced map is of order 2. Hence the degree of the composition of this map with a map $\mathbb{S}^n \rightarrow \text{Hom}(C_{2r+1}, K_{n+2})$ is zero. A very elegant and simple proof of this fact, avoiding all cohomology calculations, has recently been found by Živaljević [Živ05]. A $\mathbb{Z}_2$ -map $\mathbb{S}^n \rightarrow \mathbb{S}^n$, however, has odd degree.

For odd n , Babson and Kozlov use the same $\mathbb{Z}_2$ -map and the map between quotient spaces $\text{Hom}(C_{2r+1}, K_{n+2})/\mathbb{Z}_2 \rightarrow \mathbb{R}P^n$ it induces and show that the image of the generator of $H^n(\mathbb{R}P^n; \mathbb{Z}_2)$ in $H^n(\text{Hom}(C_{2r+1}, K_{n+2})/\mathbb{Z}_2; \mathbb{Z}_2)$ is zero. Thus for odd n they proved the following:

1.3. Theorem. *Let $r \geq 1$, $n \geq 0$. Then $w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = 0$.*

Here w_1 denotes the first Stiefel-Whitney class of the free $\mathbb{Z}_2$ -space $\text{Hom}(C_{2r+1}, K_{n+2})$ (which can be regarded as a principal $\mathbb{Z}_2$ -bundle over $\text{Hom}(C_{2r+1}, K_{n+2})/\mathbb{Z}_2$, which determines a real line bundle since $\mathbb{Z}_2 \cong O(1)$). The statement $w_1^0(X) = 0$ is to be read as $H^0(X/\mathbb{Z}_2; \mathbb{Z}_2) = 0$.

Babson and Kozlov have conjectured that Theorem 1.3 holds for all n [BK06b, Conj. 2.5], and the objective of this article is to prove this, by an argument that is valid for all n .

From Theorem 1.3 using the naturality of w_1 with respect to $\mathbb{Z}_2$ -maps and $w_1^n(\mathbb{S}^n) \neq 0$ one immediately deduces Theorem 1.1. Using only the naturality of w_1 we can also state the following stronger version which has been announced by Babson and Kozlov in [BK03].

1.4. Theorem. *Let G be a graph, $r \geq 1$ and $n \geq 0$. If $w_1^n(\text{Hom}(C_{2r+1}, G)) \neq 0$, then G is not $(n+2)$ -colourable.*

This version is not only stronger, it also stresses the importance of the $\mathbb{Z}_2$ -action and may be better suited to actual computations.

Outline of the proof of Theorem 1.3. We will construct an equivariant map of free $\mathbb{Z}_2$ -spaces $f: \text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow X_{n,r}$ where $X_{n,r}$ is a product of r spheres of dimension n with a certain operation of $\mathbb{Z}_2$. The map f misses a certain $\mathbb{Z}_2$ -invariant subspace $A_{n,r} \subset X_{n,r}$. An elementary computation shows that the n -th power of the Stiefel-Whitney class w_1 of $X_{n,r} \setminus A_{n,r}$, and hence of $\text{Hom}(C_{2r+1}, K_{n+2})$, is zero.

Subsequent developments. The proof presented here was the first proof of Theorem 1.3 and the first simple proof of Theorem 1.1. Later on, in [Sch06] the more general result that $w_1^n(\text{Hom}(C_{2r+1}, G)) = 0$ holds for all graphs G with $w_1^{n+1}(\text{Hom}(K_2, G)) = 0$ has been proven. In [Koz06b] a proof of Theorem 1.3 is given that proceeds similarly to the proof in this work but uses calculations at the cochain level instead of our calculations in the homology groups of products of spheres.

Acknowledgements. The construction presented here has been inspired by a conjecture of Péter Csorba [Cso05, Conj. 4.8] that I learned about from Frank Lutz and that has lead to their work [CL05] on Hom-complexes that are manifolds. That conjecture states that $\text{Hom}(C_5, K_{n+2})$ is homeomorphic to the Stiefel manifold $\{(x, y) \in \mathbb{S}^n \times \mathbb{S}^n : \langle x, y \rangle = 0\}$ and has been proved in [Sch05].

2. THE MAP $\text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow X_{n,r} \setminus A_{n,r}$

Let $r \geq 1$, $n \geq 0$.

2.1. Notation. We denote the vertices of C_{2r+1} by $v_0, \dots, v_{2r}$, in some cyclic order. The barycentric subdivision of the complex $\text{Hom}(C_{2r+1}, K_{n+2})$ can be described as the order complex of the poset

$$\begin{aligned} \{\phi: V(C_{2r+1}) \rightarrow \mathcal{P}(V(K_{n+2})) \setminus \{\emptyset\} : \\ \phi(v_i) \cap \phi(v_{i+1}) = \emptyset, 0 \leq i < 2r, \phi(v_0) \cap \phi(v_{2r}) = \emptyset\} \end{aligned}$$

ordered by $\phi \leq \phi' \iff \phi(v_i) \subset \phi'(v_i)$ for all i . A vertex of this subdivision can be regarded as a multi-colouring of C_{2r+1} , in which each vertex is assigned a non-empty set of colours, i.e. vertices of K_{n+2} , and adjacent vertices are assigned disjoint sets. We will always work with this subdivision. The $\mathbb{Z}_2$ -operation on $\text{Hom}(C_{2r+1}, K_{n+2})$ induced by the automorphism of C_{2r+1} that flips the edge $\{v_0, v_{2r}\}$ can be described explicitly by $(\tau\phi)(v_i) = \phi(v_{2r-i})$, where τ denotes the generator of $\mathbb{Z}_2$.

We will compare $\text{Hom}(C_{2r+1}, K_{n+2})$ to the following space.

2.2. Definition. We define the space $X_{r,n}$ by

$$X_{r,n} := \prod_{i=0}^{2r-1} \mathbb{S}^n = (\mathbb{S}^n)^{2r}$$

and a $\mathbb{Z}_2$ -operation on it by

$$\begin{aligned} (\tau x)_0 &:= -x_0, \\ (\tau x)_i &:= x_{2r-i}, \quad 0 < i < 2r. \end{aligned}$$

We will define an equivariant map $f = (f_i)_i: \text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow X_{n,r}$ as follows. The inclusion of K_2 to the edge $\{v_0, v_{2r}\}$ of C_{2r+1} induces a map

$$\text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow \text{Hom}(K_2, K_{n+2}) \approx \mathbb{S}^n.$$

This will be f_0 . The inclusion of the vertex v_i induces a map

$$\text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow \text{Hom}(K_1, K_{n+2}) \approx \mathbb{D}^{n+1}.$$

Since v_{2i} is not isolated, the image of this map is contained in (actually equal to) $\partial \mathbb{D}^{n+1} = \mathbb{S}^n$. This will be f_i for $i > 0$. We now carry out this construction in more detail.

We identify the vertices of K_{n+2} with the vertices of the $(n+1)$ -simplex Δ^{n+1} . The barycentric subdivision of the boundary of this simplex, $\text{bsd}(\partial \Delta^{n+1})$, is the order complex of the set $\{A: A \subset V(K_{n+2}), A \neq \emptyset, \mathcal{C}A \neq \emptyset\}$ ordered by inclusion. The map sending a subset of $V(K_{n+2})$ to its complement is a simplicial map of this complex to itself, and we choose a triangulation $t: |\text{bsd}(\partial \Delta^{n+1})| \rightarrow \mathbb{S}^n$ of the n -sphere that transports this map to the antipodal map of the sphere.

2.3. Definition. For $0 < i < 2r$ we define $f_i: \text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow \mathbb{S}^n$ as the composition of the simplicial map

$$\begin{aligned} \text{Hom}(C_{2r+1}, K_{n+2}) &\rightarrow \text{bsd}(\partial \Delta^{n+1}), \\ \phi &\mapsto \phi(v_i) \end{aligned}$$

with the triangulation t .

2.4. Lemma. For $0 < i < 2r - 1$ and any point $y \in \text{Hom}(C_{2r+1}, K_{n+2})$ we have $f_i(y) \neq f_{i+1}(y)$.

Proof. Let $y \in \text{Hom}(C_{2r+1}, K_{n+2})$. The point y is contained in a simplex defined by a chain $\phi_0 < \dots < \phi_s$. The points $f_i(y)$ and $f_{i+1}(y)$ are contained in the simplices defined by the chains $\phi_0(v_i) \subset \dots \subset \phi_s(v_i)$ and $\phi_0(v_{i+1}) \subset \dots \subset \phi_s(v_{i+1})$ respectively. Since $\phi_s(v_i) \cap \phi_s(v_{i+1}) = \emptyset$, these simplices have no vertex in common and hence an empty intersection. $\square$

For a poset P we can consider its corresponding *interval poset* $\{(p, q): p \leq q\}$ ordered by $(p, q) \leq (p', q') \iff (p \leq p') \wedge (q \geq q')$, the order complex of which is the *edge subdivision* $\text{esd}(\Delta(P))$ of the order complex $\Delta(P)$ of P . It has as vertices the barycentres of 0-simplices and 1-simplices of $\Delta(P)$, see [Wal88]. To see that this is actually a subdivision, note that the edge subdivision of the simplex spanned by $p_0 < \dots < p_s$ can recursively be defined as the join of the barycentre of $\langle p_0, p_s \rangle$ with the union of the edge subdivisions of $\langle p_0, \dots, p_{s-1} \rangle$ and $\langle p_1, \dots, p_s \rangle$, which agree on $\langle p_1, \dots, p_{s-1} \rangle$.

The triangulation t induces a triangulation $t': \text{esd}(\text{bsd}(\partial \Delta^{n+1})) \rightarrow \mathbb{S}^n$.

2.5. Remark. The barycentric subdivision of $\text{Hom}(K_2, K_{n+2})$ is the order complex of $\{(A, B): A, B \subset V(K_{n+2}), A \cap B = \emptyset, A, B \neq \emptyset\}$. Via $(A, B) \mapsto (A, \mathcal{C}B)$ this is isomorphic to $\text{esd}(\text{bsd}(\partial \Delta^{n+1}))$. Using t' we have a homeomorphism $\text{Hom}(K_2, K_{n+2}) \approx \mathbb{S}^n$. This homeomorphism transports the automorphism of $\text{Hom}(K_2, K_{n+2})$ given by $(A, B) \mapsto (B, A)$ to the antipodal map, also see the calculation in the proof of Proposition 2.9 below. With this fact the introduction contains a complete, albeit sketchy, proof of Theorem 1.2.

2.6. Definition. We define $f_0: \text{Hom}(C_{2r+1}, K_{n+2}) \rightarrow \mathbb{S}^n$ as the composition of the simplicial map

$$\begin{aligned} \text{Hom}(C_{2r+1}, K_{n+2}) &\rightarrow \text{esd}(\text{bsd}(\Delta^{n+1})), \\ \phi &\mapsto (\phi(v_0), \mathcal{C}(\phi(v_{2r}))) \end{aligned}$$

with the triangulation t' .

2.7. Lemma. For any point $y \in \text{Hom}(C_{2r+1}, K_{n+2})$ we have $f_0(y) \neq f_1(y)$.

Proof. Let $y \in \text{Hom}(C_{2r+1}, K_{n+2})$. The point y is contained in the interior of a unique simplex defined by a chain $\phi_0 < \dots < \phi_s$. The point $f_1(y)$ is in the image under t of the simplex spanned by $\phi_0(v_1) \subset \dots \subset \phi_s(v_1)$. The point $f_0(y)$ is in the image under t' of the interior of the simplex spanned by $(\phi_0(v_0), \mathcal{C}\phi_0(v_{2r})) \leq \dots \leq (\phi_s(v_0), \mathcal{C}\phi_s(v_{2r}))$ and hence in the image under t of the interior of the simplex spanned by $\phi_0(v_0) \subset \dots \subset \phi_s(v_0) \subset \mathcal{C}\phi_s(v_{2r}) \subset \dots \subset \mathcal{C}\phi_0(v_{2r})$. If $f_0(y) = f_1(y)$ then $\phi_0(v_1) = \phi_0(v_0)$ contradicting $\phi_0(v_1) \cap \phi_0(v_0) = \emptyset$. $\square$

2.8. Definition. We define $A_{r,n} \subset X_{r,n}$ by

$$\begin{aligned} A_{r,n} &:= \bigcup_{i=0}^{2r-1} A_{r,n}^i, \\ A_{r,n}^i &:= \{x \in X_{r,n} : x_i = x_{i+1}\}, \quad 0 \leq i < 2r-1, \\ A_{r,n}^{2r-1} &:= \{x \in X_{r,n} : x_{2r-1} = -x_0\}. \end{aligned}$$

Since $\tau A_{r,n}^i = A_{r,n}^{2r-1-i}$, the subset $A_{r,n}$ of $X_{r,n}$ is invariant under the operation of $\mathbb{Z}_2$.

2.9. Proposition. The above construction yields a $\mathbb{Z}_2$ -equivariant map

$$f: \text{Hom}(C_{2r+1}, K_{n+2}) \longrightarrow X_{n,r} \setminus A_{n,r}.$$

Proof. Let $y \in \text{Hom}(C_{2r+1}, K_{n+2})$. We assume that y has affine coordinates $(\lambda_j)_j$ in the simplex $\langle \phi_0, \dots, \phi_s \rangle$ and write this fact as $y = \sum_j \lambda_j \phi_j$. For $i > 0$ we have

$$\begin{aligned} f_i(\tau y) &= t\left(\sum_j \lambda_j (\tau \phi_j)(v_i)\right) = t\left(\sum_j \lambda_j \phi_j(v_{2r-i})\right) \\ &= f_{2r-i}(y). \end{aligned}$$

Also,

$$\begin{aligned} f_0(\tau y) &= t'\left(\sum_j \lambda_j ((\tau \phi_j)(v_0), \mathcal{C}(\tau \phi_j)(v_{2r}))\right) = t'\left(\sum_j \lambda_j (\phi_j(v_{2r}), \mathcal{C}\phi_j(v_0))\right) \\ &= t\left(\sum_j \lambda_j \left(\frac{1}{2}\phi_j(v_{2r}) + \frac{1}{2}\mathcal{C}\phi_j(v_0)\right)\right) \\ &= -t\left(\sum_j \lambda_j \left(\frac{1}{2}\mathcal{C}\phi_j(v_{2r}) + \frac{1}{2}\phi_j(v_0)\right)\right) = -t'\left(\sum_j \lambda_j (\phi_j(v_0), \mathcal{C}\phi_j(v_{2r}))\right) \\ &= -f_0(y). \end{aligned}$$

Therefore $f(\tau y) = \tau f(y)$.

For $0 < i < 2r - 1$ we have $f(y) \notin A_{r,n}^i$ by Lemma 2.4. From Lemma 2.7 follows $f(y) \notin A_{r,n}^0$ and $f(y) = \tau f(\tau y) \notin \tau A_{r,n}^0 = A_{r,n}^{2r-1}$. Thus $f(y) \notin A_{r,n}$. $\square$

$$3. \ w_1^n(X_{r,n} \setminus A_{r,n}) = 0$$

We want to determine the class $w_1^n(X_{r,n} \setminus A_{r,n}) \in H^n((X_{r,n} \setminus A_{r,n})/\mathbb{Z}_2; \mathbb{Z}_2)$. Given any equivariant map $X_{r,n} \setminus A_{r,n} \rightarrow \mathbb{S}^n$, this class is the image of the generator $\omega \in H^n(\mathbb{R}P^n; \mathbb{Z}_2)$ under the induced map $H^*(\mathbb{R}P^n; \mathbb{Z}_2) \rightarrow H^*((X_{r,n} \setminus A_{r,n})/\mathbb{Z}_2; \mathbb{Z}_2)$. The equivariant map we will use is the projection

$$p_0: X_{r,n} = \prod_{i=0}^{2r-1} \mathbb{S}^n \rightarrow \mathbb{S}^n \\ x \mapsto x_0.$$

We note that the induced map $\bar{p}_0: X_{r,n}/\mathbb{Z}_2 \rightarrow \mathbb{R}P^n$ is a locally trivial fibration with fiber $(\mathbb{S}^n)^{2r-1}$.

Since $A_{r,n}/\mathbb{Z}_2$ is a closed subset of the $2rn$ -dimensional manifold $X_{r,n}/\mathbb{Z}_2$, we have Poincaré duality at our disposal. We notationally omit the coefficient field $\mathbb{Z}_2$ from now on and consider the commutative diagram

$$\begin{array}{ccccc} & & & & H^n(\mathbb{R}P^n) \\ & & & \swarrow \bar{p}_0^* & \downarrow \\ H^n(X_{r,n}/\mathbb{Z}_2, (X_{r,n} \setminus A_{r,n})/\mathbb{Z}_2) & \xrightarrow{k^*} & H^n(X_{r,n}/\mathbb{Z}_2) & \xrightarrow{j^*} & H^n((X_{r,n} \setminus A_{r,n})/\mathbb{Z}_2) \\ \downarrow \cong & & \downarrow \cong & & \\ H_{(2r-1)n}(A_{r,n}/\mathbb{Z}_2) & \longrightarrow & H_{(2r-1)n}(X_{r,n}/\mathbb{Z}_2) & & \\ \uparrow & & \uparrow \pi_* & & \\ H_{(2r-1)n}(A_{r,n}) & \xrightarrow{l_*} & H_{(2r-1)n}(X_{r,n}) & & \end{array}$$

with j , k , and l inclusions and the isomorphisms given by Poincaré duality.

Now $w_1^n(X_{r,n} \setminus A_{r,n}) = (\bar{p}_0 \circ j)^*(\omega)$, where ω is the generator of $H^n(\mathbb{R}P)$. The Poincaré dual of $\bar{p}_0^*(\omega)$ is $\bar{p}_{0!}(\ast)$ with $\ast$ being the class of a point, i.e. the generator of $H_0(\mathbb{R}P^n)$, the Poincaré dual of ω . $\bar{p}_{0!}(\ast)$ is the orientation class (a bit of a misnomer when working with coefficients in $\mathbb{Z}_2$) of a fibre of the map $\bar{p}_0$, and therefore $\bar{p}_{0!}(\ast) = \pi_*(c_0)$ with the following notation.

3.1. Notation. For $0 \leq i < 2r$ we define $c_i \in H_{(2r-1)n}(X_{r,n})$ by

$$c_i := \underbrace{[S^n] \times \cdots \times [S^n]}_{i \text{ factors}} \times \ast \times \underbrace{[S^n] \times \cdots \times [S^n]}_{2r-i-1 \text{ factors}}.$$

3.2. Lemma. For $0 \leq i < 2r - 1$ we have $c_i + c_{i+1} \in \text{im } l_*$.

Proof. Considering the map $\Delta: \mathbb{S}^n \rightarrow \mathbb{S}^n \times \mathbb{S}^n$ given by $\Delta(x) = (x, x)$ we have $\Delta_*(c) = * \times c + c \times *$ for $c \in H_n(\mathbb{S}^n; \mathbb{Z}_2)$. With

$$\begin{aligned} \Delta_i: (\mathbb{S}^n)^{2r-1} &\rightarrow A_{r,n}^i \subset A_{r,n} \\ \Delta_i &:= \underbrace{\text{id} \times \cdots \times \text{id}}_{i \text{ factors}} \times \Delta \times \underbrace{\text{id} \times \cdots \times \text{id}}_{2r-i-1 \text{ factors}} \end{aligned}$$

we therefore have $c_i = l_*((\Delta_i)_*([\mathbb{S}^n] \times \cdots \times [\mathbb{S}^n]))$. $\square$

3.3. Lemma. *We have $\pi_*(c_r) = 0$.*

Proof. Singling out the sphere at position r of $X_{r,n}$ we obtain a $\mathbb{Z}_2$ -homeomorphism $X_{r,n} \approx Y \times \mathbb{S}^n$, where Y is a product of $2r-1$ spheres with the appropriate free $\mathbb{Z}_2$ -action and the operation on $\mathbb{S}^n$ is trivial. Under this homeomorphism, c_r corresponds to $[Y] \times *$, where $[Y]$ is the orientation class of Y . Since the quotient map $Y \rightarrow Y/\mathbb{Z}_2$ is of even degree, the image of $[Y] \times *$ in $H_{(2r-1)n}((Y \times \mathbb{S}^n)/\mathbb{Z}_2) = H_{(2r-1)n}((Y/\mathbb{Z}_2) \times \mathbb{S}^n)$ is zero. $\square$

3.4. Proposition. *We have $w_1^n(X_{r,n} \setminus A_{r,n}) = 0$.*

Proof. As noted before, $w_1^n(X_{r,n} \setminus A_{r,n}) = j^*(\bar{p}_0^*(\omega))$ with $\bar{p}_0^*(\omega)$ Poincaré dual to $\pi_*(c_0)$. By the preceding lemmas

$$\pi_*(c_0) = \pi_*(c_0 + c_r) = \pi_*\left(\sum_{i=0}^{r-1} (c_i + c_{i+1})\right) \in \text{im}(\pi_* \circ l_*).$$

Using the commutative diagram it follows that $\bar{p}_0^*(\omega) \in \text{im } k^* = \ker j^*$. $\square$

4. RESULT

Our main result appears as Theorem 1.3 in the introduction.

Theorem. *Let $r \geq 1$ and $n \geq 0$. Then $w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = 0$.*

Proof. This follows directly from Proposition 3.4, since w_1 is characteristic: With the map $\bar{f}: \text{Hom}(C_{2r+1}, K_{n+2})/\mathbb{Z}_2 \rightarrow (X_{r,n} \setminus A_{r,n})/\mathbb{Z}_2$ induced by the equivariant map f from Proposition 2.9 we have

$$w_1^n(\text{Hom}(C_{2r+1}, K_{n+2})) = \bar{f}^*(w_1^n(X_{r,n} \setminus A_{r,n})) = \bar{f}^*(0) = 0$$

as claimed. $\square$

REFERENCES

- [BK03] BABSON, E. and KOZLOV, D. N. Topological obstructions to graph colorings. *Electron. Res. Announc. Amer. Math. Soc.*, **9**:61–68, 2003.
- [BK06a] —. Complexes of graph homomorphisms. *Isr. J. Math.*, **152**:285–312, 2006. math.CO/0310056.
- [BK06b] —. Proof of the Lovász conjecture. *Annals of Mathematics*, 2006. In press, math.CO/0402395.
- [CL05] CSORBA, P. and LUTZ, F. H. Graph coloring manifolds, 2005. 22 pp., math.CO/0510177.
- [Cso05] CSORBA, P. *Non-tidy Spaces and Graph Colorings*. Ph.D. thesis, ETH Zürich, 2005.
- [Koz06a] KOZLOV, D. N. Chromatic numbers, morphism complexes, and Stiefel-Whitney characteristic classes. In MILLER, E., REINER, V., and STURMFELS, B., eds., *Geometric Combinatorics*. American Mathematical Society, 2006. In press, math.AT/0505563.

- [Koz06b] —. Cobounding odd cycle colorings, 2006. Preprint, math.AT/0602561.
- [Lov78] LOVÁSZ, L. Kneser's conjecture, chromatic number and homotopy. *J. Combinatorial Theory, Ser. A*, **25**:319–324, 1978.
- [Sch05] SCHULTZ, C. Small models of graph colouring manifolds and the Stiefel manifolds $\text{Hom}(C_5, K_n)$, 2005. Preprint, 19 pp., math.CO/0510535.
- [Sch06] —. Graph colourings, spaces of edges and spaces of circuits, 2006. Preprint, math.CO/0606763.
- [Wal88] WALKER, J. W. Canonical homeomorphisms of posets. *European J. Combin.*, **9**(2):97–107, 1988. ISSN 0195-6698.
- [Živ05] ŽIVALJEVIĆ, R. T. Combinatorial groupoids, cubical complexes, and the Lovász conjecture, 2005. 28 pp., math.CO/0510204.

INSTITUT FÜR MATHEMATIK, MA 6-2, TECHNISCHE UNIVERSITÄT BERLIN, D-10623 BERLIN, GERMANY

E-mail address: `carsten@codimi.de`